



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

going to breed. There can be no reasonable doubt that this species breeds in northern New Jersey.—W. DEW. MILLER AND LUDLOW GRISCOM, *American Museum of Natural History, New York City.*

Hermit Thrush's Nest in Unusual Location.—At Jefferson Highland N. H., the Hermit Thrush (*Hylocichla guttata pallasi*) is a common summer resident. In the season of 1919, six singing males have been within hearing of my home, outnumbering the Robin two to one. And their disposition to sing freely at all hours of the day from an hour before sunrise to a half-hour after sunset, far surpasses that of our local Robins. One pair of Hermits has been located at the wood border below our garden and has been frequently seen in the garden and nearby orchard. When the season of ripe blackberries had come and I was gathering berries on August 9 in the plot of considerable size within our garden, I came upon a nest lodged on several of the canes within about a foot of their tips and *four feet* or so *above the ground*. One fledgling about ready to fly was in the nest, and I surmised that two or three others had probably already flown. The mother bird came and perched on the top of a bean pole standing, perhaps, thirty feet from her nest and showed no excited anxiety over my presence, continuing to hold her perch for some time and quietly giving her hissing call only. The next day when the blackberry plot was visited, the fledgling left the nest upon my approach. This nest rested firmly on several canes and was concealed from casual view by the leaves thickly surrounding it; while clusters of berries hung all about it, so it had not been discovered until on the day named the branches were drawn aside a little in gathering the fruit. The location in the plot of blackberries was on the outer edge southward, the plot having a width of about twenty feet, and was one hundred and fifty feet from the border of the woods, thus well up therefrom in the garden. The male bird continued singing up to August 16 inclusive; on that day I heard him sing a few times at 6 A. M. and again at 12 M., as he had done the preceding day. But this was the last voicing of his beautiful song for the season. Subsequently, an occasional call only was heard, and like his brother Hermits in the neighborhood he was for the most part silent, without even expressing himself in either of his four distinctly different call-notes.—HORACE W. WRIGHT, *107 Pinckney St., Boston, Mass.*

Peculiar Nesting of Hermit Thrushes.—How far the nesting of birds may be influenced by friendly contact with man is worthy of careful study and observation by all bird lovers. The peculiar nesting plan of a pair of Hermit Thrushes, near the shore of Asquam Lake, Holderness, N. H., is of especial interest and may lead to other observations of a like peculiarity. In all our previous observations of their nesting, the nests were found among low bushes on damp ground. Mr. F. Schuyler Matthews writes that he has found them on low bushes near the ground. How far the following facts may have been influenced by association with man, can only be conjectured.